

Multi-Probiotic 4000

Friendly flora to promote intestinal balance

DOUGLAS LABORATORIES®

Product Data

Description

Each Multi-Probiotic 4000 capsule, provided by Douglas Laboratories, contains over four billion beneficial organisms, including representatives of both the lactobacillus and bifidobacterium genera, mixed in a base of prebiotic fructooligosaccharide.

Functions

Oral supplements of live, beneficial intestinal microorganisms for nutritional health and well-being are known as probiotics. Probiotic dietary lactobacilli colonize the intestinal tract, depending on the specific bacterium-to-host affinity. Bacteria are continually in competition for colonization space. The original bacterial colonies have an advantage over transient bacteria. Nevertheless, the composition of the intestinal microflora is dynamic and constantly changing.

If harmful microorganisms proliferate, the equilibrium is disturbed, and it becomes difficult even for indigenous

bacteria to maintain their necessary territory to colonize the intestine.

Antibiotics tend to kill off both beneficial and harmful bacteria, and thus may also disturb the normal, healthy balance of intestinal microorganisms.

When potent probiotic supplements, such as Multi-Probiotic 4000, are supplied regularly, chances are greatly increased for establishing and maintaining a healthy population of beneficial intestinal microorganisms. Once established, a normal intestinal microflora rich in lactobacilli creates acidic conditions that are unfavorable for the settlement of pathogenic microorganisms.





Indications

Multi-Probiotic 4000 capsules may be a useful dietary supplement for those who wish to support their intestinal microflora with meaningful amounts of beneficial microorganisms.

Formular (#7497)

Each Multi-Probiotic 4000 capsule contains over 4 billion (4000 x 1 million) organisms in the following potencies:

Lactobacillus. acidophilus (DDS-1™).....	1.15 Billion
L. Rhamnosus.....	1.15 Billion
L. Rhamnosus (Type B, Bifidus).....	775 Million
Bifidobacterium lactis.....	275 Million
Bifidobacterium longum.....	275 Million
B. Bifidum.....	275 Million
Streptococcus thermophilus.....	150 Million

In a base of FOS (fructooligosaccharides), special carbohydrates, enzymes and ulmus fulva

Suggested Use

Adults take 1 capsule daily or as directed by physician.

Side Effects

No adverse effects have been reported.

Storage

Store in a cool, dry place, away from direct light. Keep out of reach of children.

References

Cebra JJ. Influences of microbiota on intestinal immune system development. Am J Clin Nutr 1999;69:1046S-1051S.

Collins MD, Gibson GR. Probiotics, prebiotics, and synbiotics: approaches for modulating the microbial ecology of the gut. Am J Clin Nutr 1999;69:1052S-1057S.

Donnet-Hughes A, Rochat F, Serrant P, Aeschlimann JM, Schiffrin EJ. Modulation of nonspecific mechanisms of defense by lactic acid bacteria: effective dose. J Dairy Sci 1999;82:863-9.

Elmer GW, Surawicz CM, McFarland LV. Biotherapeutic agents. A neglected modality for the treatment and prevention of selected intestinal and vaginal infections [see comments]. Jama 1996;275:870-6.

Gibson GR. Dietary modulation of the human gut microflora using prebiotics. Br J Nutr 1998;80:S209-12.

Goldin BR. Health benefits of probiotics. Br J Nutr 1998;80:S203-7.

Jacobsen CN, Rosenfeldt Nielsen V, Hayford AE, Moller PL, Michaelsen KF, Paerregaard A, Sandstrom B, Tvede M, Jakobsen M. Screening of probiotic activities of forty-seven strains of lactobacillus spp. by In vitro techniques and evaluation of the colonization ability of five selected strains in humans [In Process Citation]. Appl Environ Microbiol 1999;65:4949-56.

Kirjavainen PV, Ouwehand AC, Isolauri E, Salminen SJ. The ability of probiotic bacteria to bind to human intestinal mucus.



santē | chiropractic
wellness

1-1224 Prom. Place d'Orleans Dr.
Ottawa ON K1C 7K3
613-837-2883 | santechiropractic.com

FEMS Microbiol Lett 1998;167:185-9.

Pessi T, Sutas Y, Saxelin M, Kallioinen H, Isolauri E. Antiproliferative effects of homogenates derived from five strains of candidate probiotic bacteria [In Process Citation]. Appl Environ Microbiol 1999;65:4725-8.

Tejada-Simon MV, Lee JH, Ustunol Z, Pestka JJ. Ingestion of yogurt containing Lactobacillus acidophilus and Bifidobacterium to potentiate immunoglobulin A responses to cholera toxin in mice. J Dairy Sci 1999;82:649-60.

Venturi A, Gionchetti P, Rizzello F, Johansson R, Zucconi E, Brigidi P, Matteuzzi D, Campieri M. Impact on the composition of the faecal flora by a new probiotic preparation: preliminary data on maintenance treatment of patients with ulcerative colitis. Aliment Pharmacol Ther 1999;13:1103-8.

Wang X, Ma G, Zheng B, Tian H. [Effects of SL-probiotic preparation on the body weight and phagocytosis of white mice]. Wei Sheng Wu Hsueh Pao 1995;35:455-9.

† These statements have not been evaluated by the Food and Drug Administration.

This product is not intended to diagnose, treat, cure, or prevent any disease.